

# **GREENSIEW** Fairway Formula Spring Fertilizer with Crabgrass Preventer

Prevents crabgrass for up to 120 days

Extended time release feeding of nitrogen greens into summer without excess growth

ACTIVE INGREDIENT: Dithiopyr..... OTHER INGREDIENTS TOTAL

... 0.13% . <u>99.87%</u> 100.00%

CAS. No. 97886-45-8 EPA Reg. No. 961-392

(L) (D) EPA Est. No. 961-PA-01, 961-IL-01 Superscript used is first letter of lot number.

## KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for First Aid and additional Precautionary Statements READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS. If terms are not acceptable, return the unopened container at once.

In the State of New York, do not exceed 4.4 lbs. product per 1000 sq. ft. (0.25 lb. active ingredient per acre) per year. Do not apply more than once per year to the same area.

## **Covers 5,000 sq. ft. 26–0–5** Net Weight 17 lbs. (7.7 kg.)

## Prevents crabgrass for up to 120 days

## Extended time release feeding of nitrogen greens

## into summer without excess growth

#### 

CAS. No. 97886-45-8 EPA Reg. No. 961-392 (L) (D) EPA Est. No. 961-PA-01, 961-IL-01 Superscript used is first letter of lot number.

### KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for First Aid and additional Precautionary Statements READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

If terms are not acceptable, return the unopened container at once.

In the State of New York, do not exceed 4.4 lbs. product per 1000 sq. ft. (0.25 lb. active ingredient per acre) per year. Do not apply more than once per year to the same area.

## **Covers 5,000 sq. ft. 26–0–5** Net Weight 17 lbs. (7.7 kg.)

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

CAUTION: Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

If in eyes:       • Hold eye open and rinse slowly and gently with water for 15-20 minutes.         • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.         • Call a poison control center or doctor for treatment advice.         If inhaled:       • Move person to fresh air.         • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.         • Call a noison control center or doctor for further treatment advice.	First Aid			
<ul> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.</li> </ul>	If in eyes:	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.		
	If inhaled:	If person is not breathing, call 911 or an ambulance, then give artificial respiration.		

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of an emergency involving this product, call day or night (888) 208-1368.

## **PRODUCT INFORMATION FOR TURF USE**

#### **Directions for Use**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This product contains Dimension", a herbicide that can be applied up to 4 weeks later than other crabgrass preventers and still provides season-long control. Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer prevents crabgrass and other listed grass and broadleaf weeds up to 4 months, when applied prior to their germination in lawns and ornamental turfgrasses. It also provides postemergence control of seedling crabgrass for up to 4 weeks after it germinates.

Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer should be applied to lawns and ornamental turfgrasses that are well established (See "Precautions and Restrictions" section below for a list of the tolerant turfgrasses). It will not harm nearby established ornamentals when used according to directions, given below.

#### **Precautions and Restrictions**

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES. NOT FOR USE ON NEWLY SEEDED LAWNS.

Use of Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Do not apply to lawns or turfgrasses that are under stress.

## **APPLICATION DIRECTIONS**

Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer can be applied to the following turfgrasses when used as directed:

**Cool-Season Turfgrasses -** Creeping Bentgrass, Fine Fescues, Kentucky Bluegrass, Perennial Ryegrass, Tall Fescues.

Warm-Season Turfgrasses - Bahiagrass, Bermudagrass, Buffalograss, Carpetgrass, Centipedegrass, St. Augustinegrass, Zoysiagrass.

#### **Application Timing**

Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer is to be applied in the spring or in the fall prior to weed seed germination for control of listed weeds in established lawns and ornamental turgrasses. When this product is applied in the fall, a sequential or second application must be made the following spring to provide season-long control of weeds. If this product is applied in the fall, DO NOT reseed, overseed or sprig turgrasses before 12 weeks after application due to potential turgrass injury to the newly reseeded, overseeded or sprigged turgrasses.

When establishing a new lawn of Kentucky Bluegrass, Perennial Ryegrass and/or Tall Fescue, this product may be applied 2 weeks after the first sign of grass germination. For other turfgrass, wait until the grass is well established (2 mowings) before applying this product. Do not apply this product within an 8 week period prior to seeding, reseeding, overseeding or sprigging.

Improved weed control may occur when this product is applied within a few days after mowing and further mowing is delayed until the turfgrass has been watered or received rainfall to dissolve the fertilizer granules.

Note: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

#### Length of Control

Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer will provide up to 4 months of weed control if applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, and intensity of weed pressure.

#### Post Emergence Crabgrass Control

For post emergent crabgrass control, apply Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer when crabgrass is small (3-4 leaf). It will not control large tillered crabgrass.

#### Environmental Hazards:

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather will help to ensure that wind does not blow pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

Reseeding, overseeding, or sprigging of turfgrasses into areas that have been previously treated with this product should be delayed until after 8 weeks from the time of application. Reseeding, overseeding, or sprigging before 8 weeks or earlier after application may injure or prevent the establishment of the newly reseeded, overseeded or sprigged turfgrasses.

Do not make an application of this product later than 4 weeks after crabgrass germination. Do not expect control with applications made later than 4 weeks after crabgrass germination.

**DO NOT** apply more than 8.8 lbs. of this product per 1,000 sq. ft. per application, and no more than 26.4 lbs. of this product per 1000 sq. ft. per year.

In the State of New York, do not exceed 4.4 lbs. product per 1000 sq. ft. (0.25 lb. active ingredient per acre) per year.

In the State of Maryland apply this product at no more than 3.4 lbs. of product per 1,000 sq. ft.

**DO NOT** use clippings from treated turf for mulching, vegetables, vines, or fruit trees from which feed or food products may be harvested.

Keep people and pets off treated areas until dust has settled.

#### Weeds Controlled and/or Suppressed

**Grass Weeds -** Annual Bluegrass, Annual Ryegrass, Barnyardgrass, Crabgrass, Crowfootgrass, Goosegrass, Green Foxtail, Kikuyugrass (seedling), Perennial Ryegrass, Smutgrass (seedling), Yellow Foxtail.

**Broadleaf Weeds -** Bittercress, Buttercup Oxalis, Carolina Geranium, Chickweed, Common Lespedeza, Common Purslane, Corn Speedwell, Creeping Oxalis, Henbit, Parsley-piert, Pineappleweed, Prostrate Spurge, Spotted Spurge, Yellow Woodsorrel.

	Product Use Rate Table (Lbs. Per 1,000 Sq. Ft.)			
	2-3	3-4	Early Post	
Region	Months Control	Months Control	Emergence Control	
North	2.2	3.1	3.9 - 8.8	
South	3.4	3.9 - 4.5	3.9 - 8.8	

Use the higher rate in areas of high weed pressure



#### Application Equipment and Instructions

Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer can be applied with a drop or rotary type spreader designed to apply granular fertilizers. For best results, apply this product evenly and uniformly over the turfgrass area, avoiding streaking, skips or overlaps. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Adjust the spreader settings so the spreader delivers the appropriate application rate as specified above.

A more uniform application can be made by spreading one-half of the specified amount of this product over the turgrass area and then applying the remaining one-half at a right angle to the previous direction. Check spreader frequently to ensure it is functioning properly and distributing the granules uniformly.

requerity to ensure it is functioning property and distributing the granules uniformity.

coverage slightly, and shut off spreader before making hard turns.

Apply in a single pass at a rate of 3.4 lbs. per 1,000 sq. ft. Start on the outside

border and apply in a circular path working toward the center. Overlap the

## SPREADER SETTINGS

#### **Drop Spreader Settings**

Apply in a single pass at a rate of 3.4 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.



<b>Drop Spreader</b> Earthway Drop	<b>Setting</b> (3.4 lbs. / 1000 Sq. Ft.) 14
Republic Drop	41/2
Scotts AccuGreen Drop	4
Sta-Green	4

To calibrate your spreader: 1 lb. of material should cover approximately 294 sq. ft.

If your spreader model is not listed, go to our website www.greenviewfertilizer.com, or call toll-free 800-233-1067,

8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

## FERTILIZER GUARANTEE

#### GUARANTEED ANALYSIS

Total Nitrogen (N)         26           7.6% Ammoniacal Nitrogen         6.0% Water Insoluble Nitrogen*           1.8% Urea Nitrogen         10.6% Other Water Soluble Nitrogen*           Soluble Potash (K <sub>2</sub> O)         10           Sulfur (S)         11.0% Combined Sulfur (S)	5% 5% 0%
Derived from: Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash.         *16.6% Slowly Available Nitrogen from Methylene Ureas.         F699	

#### Storage and Disposal

**Broadcast Spreader Settings** 

Broadcast Spreader Agri-Fab Rotary

Republic EZGro Rotary

Scotts

Sta-Green

Earthway EV-N-Spred Rotary

Scotts Deluxe Edgeguard Rotary

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store product in its original container in a cool, dry, locked place out of the reach of children and animals. **PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **IF EMPTY:** Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-lebsea.com

Warranty

Note To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

#### Manufactured by:

LEBANON SEABOARD CORPORATION 1600 EAST CUMBERLAND STREET LEBANON, PA 17042

For more information on our complete line of GreenView<sup>®</sup> Lawn and Garden Products, visit our website at www.greenviewfertilizer.com

#### Greenview Yard Wide Guarantee

Greenview guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem, and send proof of purchase to Lebanon Seaboard Corporation, 1600 East Cumberland Street, Lebanon, PA 17042.

GreenView<sup>®</sup> is a registered trademark of Lebanon Seaboard Corporation



Setting (3.4 lbs. / 1000 Sq. Ft.)

Δ

15

9 4½

241/2

3

21-29822

G/19