

FEEDS YOUR LAWN UP TO **8** WEEKS

GREENVIEW®

WEED & FEED

Kills 250+ Weeds*

Thickens turf to crowd out weeds

Apply when listed weeds are growing



ACTIVE INGREDIENTS:
 2,4-D, 2-ethylhexyl ester..... 1.108%
 Mecoprop-p acid.....0.167%
 Dicamba acid.....0.071%
OTHER INGREDIENTS:..... 98.654%
TOTAL100.000%

THIS PRODUCT CONTAINS:
 14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%. CAS 1928-43-4.
 3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.167%. CAS 16484-77-8.
 1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071%. CAS 1918-00-9
 Isomer Specific by AOAC Methods
 TRIMEC is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN CAUTION
 See back panel for First Aid and additional Precautionary Statements

*See Use Directions for complete list of treated Weeds

Covers 2,500 sq ft | Net Wt 7lb (3.17 kg)

GREENVIEW®

WEED & FEED

Ideal for Smaller Lawns • Great for Spot Treatment of Any Size Lawn

STOP! READ THE ENTIRE LABEL FIRST.

OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

- If in eyes:
- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Emergency Medical Treatment Information call (888)208-1368.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Do not apply this product in a way that will contact any person or pet either directly or through drift.
- This product is for residential use.
- Do not allow people (other than applicator) or pets on treatment area during treatment. Do not enter treatment area until dust has settled.

Application Instructions

GreenView Weed & Feed contains Trimec® Herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many other listed weeds commonly found in home lawns. In addition, a 27-0-4 fertilizer provides a supply of lawn food. Possible retreatment may be necessary if (a) the lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made after 30 days. Treated areas may be seeded 3 weeks after application.

USE PRECAUTIONS AND RESTRICTIONS:

Do not use on dichondra, carpetgrass, Floratam® St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are present.

Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.

Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of or over surface roots of trees, shrubs, and other ornamentals.

Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

HOW MUCH TO APPLY:

For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass, and colonial bentgrass. Apply 2.8 pounds of product per 1,000 square feet of turfgrass. A second application may be made after 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

GreenView Weed & Feed Kills more than 250 weeds as listed!

Amaranth, purple	Clover, red	Hoary cress	Plantain, blackseed	Strawberry, India mock
Amaranth, slender	Clover, strawberry	Horsenettle	Plantain, bracted	Tansy ragwort
Aster, heath	Clover, yellow sweet	Horseweed	Plantain, broadleaf	Tanweed
Aster, white heath	Clover, white sweet	Jimsonweed	Plantain, buckhorn	Thistle, blessed
Aster, purple	Clover, white	Johnny-Jumpup violet	Plantain, hoary	Thistle, bull
Aster, white prairie	Cockle	Knave	Plantain, narrowleaf	Thistle, canada
Artichoke	Cocklebur	Knotweed, prostrate	Plantain, slender	Thistle, fiodman(prairie)
Austrian fieldcress	Cocklebur, oriental	Kochia	Plantain, woolly	Thistle, musk
Bedstraw, Catchweed	Cocklebur, spiny	Lambsquarters	Poison ivy	Thistle, platte
Bedstraw, smooth	Common mullein	Lespedeza	Poison oak	Thistle, plumless
Bedstraw, Northern	Corn Spurry	Mallow, alkali	Pokeweed	Thistle, russian
Bedstraw, pineywoods	Creeping Beggarweed	Mallow, bristly	Prairie Sunflower	Thistle, scotch
Beggarticks, hairy	Creeping Jenny	Mallow, common	Prickly lettuce	Thistle, wavyleaf
Betony, Florida	(creeping charlie)	Mallow, dwarf	(compass plant)	Thistle, yellow (horrible)
Bindweed, Field	Cupid shaving Brush	Mallow, venice	Prickly sida	Trailing crownvetch
Bindweed, Hedge	Cudweed	Marcela	Prostrate knotweed	Velvetleaf
Bitter Wintercress	Daisy, English	Matchweed	Puncturevine	Vervain, blue
Bittercress, hairy	Daisy, oxeye	Mexicanweed	Purslane, common	Vervain, hoary
Bitterweed	Dandelion, common	Moneywort	Pusley, brazil	Vervain, prostrate
Black-eyed Susan	Dandelion, false	Morningglory, bigroot	Pusley, florida	Vervain, tall
Black Medic	Day flower	Morningglory, ivyleaf	Pusley, large flower	Vetch, bird
Blood flower milkweed	Deadnettle	Morningglory, mexican	Ragweed, bur	Vetch, hairy
Blue lettuce	Dichondra	Morningglory, red	Ragweed, common	Vetch, milk
Brassbuttons	Dock, broadleaf	Morningglory, tall	Ragweed, lanceleaf	Vetch, narrowleaf
Bristly oxtongue	Dock, curly	Morningglory, wolly	Ragweed, western	Vetch, wild
Broomweed	Dogbane	Mugwort	Redstem filaree	Virginia creeper
Buckhorn	Dogfennel	Mustard, ball	Rough cinquefoil	Western clematis
Bullnettle	Dollarweed (Pennywort)	Mustard, black	Scarlet pimpernel	Western salsify
Burclover	Dove weed	Mustard, blue	Shepardspurse	Wild aster
Burdock	Elderberry	Mustard, hedge	Smartweed, ladysthumb	Wild buckwheat
Burweed, lawn	Falseflax	Mustard, Indian	Smartweed, pale	Wild carrot
(spurweed)	False sunflower	Mustard, tansy	Smartweed, pennsylvania	Wild four-o'clock
Buttercup, bulbous	Fiddleneck	Mustard, tumble	Smooth chaff-flower	Wild garlic
Buttercup, creeping	Filaree, redstem	Mustard, white	Smoothdock	Wild lettuce
Buttonweed, common	Filaree, whitestem	Mustard, wormseed	Sorrel, red (sheep)	Wild marigold
(poorjoe)	Fleabane, daisy (annual)	Nettle, stinging (burning)	Southistle, annual	Wild mustard
Buttonweed, virginia	Fleabane, rough	Nettle, tall	Southistle, spiny	(yellow rocket)
Carolina geranium (wild)	Galinsoga, hairy	Nettle, wood	Spanish needles	Wild onion
Carpetweed	Galinsoga, smallflower	Parsley-piert	Speedwell, bi-lobed	Wild pansnip
Catnip	Goldenrod	Parsnip	Speedwell, common	Wild radish
Catsear, spotted	Ground ivy	Pearlwort	Speedwell, corn	Wild rape
(false dandelion)	Groundsel	Pennycress (frenchweed)	Speedwell, ivyleaf	Wild strawberry
Chamber bitter	Gumweed	Pepperweed, perennial	Speedwell, persian	Wild sweet potato
Chickweed, common	Hairy fleabane	Pepperweed, virginia	Speedwell, purslane	Wild violet
Chickweed, mouseear	Hawkweed, meadow	(Peppergrass)	Speedwell, slender	Woodsorrel, creeping
Chickweed, sticky	Hawkweed, mouseear	Pigweed, prostrate	Speedwell, snow	Woodsorrel, violet
Chicory	Hawkweed, orange	Pigweed, redroot	Pigweed, thymeleaf	Woodsorrel, yellow
Cinquefoil	Hawkweed, yellow	Pigweed, russian	Spiny amaranth	(Oxalis)
Clover, crimson	Healall	Pigweed, smooth	Spurge, spotted	Woolly croton
Clover, alyce	Heartleaf drymary	Pigweed, tumble	Spurge, prostrate	Wormseed
Clover, hop	Hemp	Pineapple weed	Star of Bethlehem	Yarrow, common
Clover, rabbit foot	Henbit	Plains coreopsis (tickseed)		Yarrow, western
				Yarrow, yellow

Maximum use rate statement

Do not apply more than 2.8 pounds of product per 1000 square feet of turfgrass per application. If necessary, a second application may be made no sooner than 30 days after the first. Do not apply more than 5.6 pounds of product per 1000 square feet per year.

The maximum broadcast application rate per acre is 1.28 lb 2,4-D ae, 0.29 lb MCPP-p ae and 0.12 lb dicamba ae per acre per application. The maximum seasonal rate is 2.56 lb 2,4-D ae, 0.58 lb MCPP-p ae and 0.25 lb dicamba ae per acre.

WHEN TO APPLY:

Apply when listed weeds are growing.


Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.

APPLICATION INSTRUCTIONS:

1. Mow lawn to normal height, 1 to 2 days before application.
2. For best results lawn must be MOIST and the air calm, when product is applied. When the lawn is moist, the product granules stick better to the leafy parts of the weeds, ensuring that the weed-killing ingredients are absorbed by the weed plant.
 - TIP: Apply after a heavy dew or rain, or on a thoroughly watered lawn.
 - TIP: DO NOT apply to a moist lawn IF rain is expected within 24 hours of application.
 - TIP: DO NOT water in immediately after application as the granules need to stick to the weed plant leaf for 1-2 days to be most effective.
3. Apply when weeds are young and actively growing.
4. Avoid applying when temperatures are consistently above 90°F.
5. After a 1-2 day period after application, if it has not rained - the lawn should be thoroughly watered.

Spreader Directions:



Determining Your Lawn's Square Footage



To determine how much of this product to apply, you must determine the square footage of your lawn. An easy way to do this is to walk the length of your lawn, figuring that one pace equals about 3 feet. Do the same with the width of the lawn. Then multiply the length by the width to arrive at the total. Make sure you subtract the square footage of your home and driveway when calculating the total square footage of your lawn. If your lawn has an irregular shape, try dividing it up into a number of square areas. Calculate the square footage for each area, then add the figures together to reach your total lawn size.

Broadcast Spreader	Setting	Drop Spreader	Setting	Handheld Spreader	Setting
	2.8 Lbs./1,000 Sq. Ft.		2.8 Lbs./1,000 Sq. Ft.	2.8 Lbs./1,000 Sq. Ft.	
Agri-Fab Rotary	4	Earthway	14	Scotts Wizz	3½
Earthway Rotary	14	Republic Drop	4	Scotts Whirl	3½
Republic Rotary	6½	Scotts	3½	Earthway EVN-Spred	3
Scotts Deluxe Edgeguard	4	Sta-Green Drop (1-5)	3½		
Scotts EasyGreen	23				
Sta-Green Rotary (1-10)	3				

Broadcast Spreader: Apply in a single pass by starting on the outside border and apply in a circular path working toward the center.



Drop Spreader: First make two passes along the top of the area you'll be covering. Then make two passes along the bottom area. Then move back and forth along the long part of the lawn. On each pass, as you reach the strips at the top and bottom of the area to be covered, shut off the spreader. It's also important to shut off the spreader when you are stopped.

Note: if spreader model does not appear above please call 800-233-1067, 8 am - 5 pm ET, Monday - Friday for additional model settings. The above spreader settings are approximate. To check your spreader: 1 lb. of material should cover approximately 385 sq. ft. (2.8 lbs. of material per 1,000 sq. ft.)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

Pesticide Disposal and Container Handling: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Questions or Comments? Call 1-800-233-1067.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

27-0-4

GUARANTEED ANALYSIS

Total Nitrogen (N)	27%
3.4% Ammoniacal Nitrogen	
2.2% Water Insoluble Nitrogen*	
18.0% Urea Nitrogen	
3.4% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	4%
Sulfur (S)	3.8%
3.8% Combined Sulfur (S)	
Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.	
* 5.6% Slowly Available Nitrogen from Methylene Ureas.	F699

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-lebsea.com>



Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem and simply send proof of purchase to Lebanon Seaboard Corporation, 1600 East Cumberland Street, Lebanon, PA 17042.

*For more information on our complete line of GreenView® Lawn and Garden Products, visit our website at www.greenviewfertilizer.com
*GreenView® is a registered trademark of Lebanon Seaboard Corporation.

Manufactured by:
Lebanon Seaboard Corporation
1600 East Cumberland Street
Lebanon, PA 17042

EPA Reg. No. 961-415

EPA Est. No. 961-PA-01, 961-IL-01

Superscript used is first letter of lot number.



21-29853