

GREENVIEW®

BROADLEAF WEED CONTROL PLUS LAWN FOOD



Helps lawns retain water to
protect against heat & drought



Speeds root growth to
improve nutrient absorption

Feeds your lawn up to 8 weeks

Apply when listed weeds are
growing in spring or early fall



For more information, visit www.greenviewfertilizer.com

**Kills Over 250
Weeds as listed**



NET WEIGHT: 13 lbs. (5.9 kg)
Covers 5,000 sq. ft.

ACTIVE INGREDIENTS:
2,4-D, 2-ethylhexyl ester 1.108%
Mecoprop-p acid 0.167%
Dicamba acid 0.071%
OTHER INGREDIENTS: 98.654%
TOTAL 100.000%

THIS PRODUCT CONTAINS:
14.70 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%. CAS 1928-43-4.
3.34 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.167%. CAS 16484-77-8.
1.42 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.071%. CAS 1918-00-9

Isomer Specific by AOAC Methods
TRIMEC is a registered trademark of PBI/Gordon Corporation.

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See back panel for First Aid and
additional Precautionary Statements.

GREENVIEW® BROADLEAF WEED CONTROL
PLUS LAWN FOOD

STOP! READ THE ENTIRE LABEL FIRST.

OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

IF IN EYES:

•Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.

• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

For Emergency Medical Treatment Information call (888)208-1368.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Do not apply this product in a way that will contact any person or pet either directly or through drift.
- This product is for residential use.
- Do not allow people (other than applicator) or pets on treatment area during treatment. Do not enter treatment area until dust has settled.

Application Instructions

GreenView Broadleaf Weed Control plus Lawn Food contains Trimec® Herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many other listed weeds commonly found in home lawns. In addition, a 27-0-4 fertilizer provides a supply of lawn food. Possible retreatment may be necessary if (a) the lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made after 30 days. Treated areas may be seeded 3 weeks after application.

USE PRECAUTIONS AND RESTRICTIONS:

Do not use on dichondra, carpetgrass, 'Floritam' St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are present.

Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.

Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of or over surface roots of trees, shrubs, and other ornamentals.

Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

HOW MUCH TO APPLY:

For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass , zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass, and colonial bentgrass. Apply 2.6 pounds of product per 1,000 square feet of turfgrass. A second application may be made after 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

Maximum Use Rate Statement

Do not apply more than 2.6 pounds of product per 1000 square feet of turfgrass per application. If necessary, a second application may be made no sooner than 30 days after the first. Do not apply more than 5.2 pounds of product per 1000 square feet per year.

The maximum broadcast application rate per acre is 1.28 lbs. 2,4-D ae, 0.29 lb. MCPP-p ae and 0.12 lb. dicamba ae per acre per application. The maximum seasonal rate is 2.56 lbs. 2,4-D ae, 0.58 lb. MCPP-p ae and 0.25 lb. dicamba ae per acre.

WHEN TO APPLY:

Apply when listed broadleaf weeds are actively growing in spring or early fall.

Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.

GreenView® Broadleaf Weed Control plus Lawn Food Kills more than 250 weeds as listed!

Amaranth, purple

Amaranth, slender

Aster, heath

Aster, white heath

Aster, purple

Aster, white prairie

Artichoke

Austrian fieldcress

Bedstraw, Catchweed

Bedstraw, smooth

Bedstraw, Northern

Bedstraw, pineywoods

Beggarticks, hairy

Betony, Florida

Bindweed, Field

Bindweed, Hedge

Bitter Wintercress

Bittercress, hairy

Bitterweed

Black-eyed Susan

Black Medic

Blood flower milkweed

Blue lettuce

Brassbuttons

Bristly oxtongue

Broomweed

Buckhorn

Bullnettle

Burclover

Burdock

Burweed, lawn

(spurweed)

Buttercup, bulbous

Buttercup, creeping

Buttonweed, common

(poorjoe)

Buttonweed, Virginia

Carolina geranium (wild)

Carpetweed

Catnip

Catsear, spotted

(false dandelion)

Chamber bitter

Chickweed, common

Chickweed, mouseear

Chickweed, sticky

Chicory

Cinquefoil

Clover, crimson

Clover, Alyce

Clover, hop

Clover, rabbit foot

Clover, red

Clover, strawberry

Clover, yellow sweet

Clover, white sweet

Clover, white

Cockle

Cocklebur

Cocklebur, oriental

Cocklebur, spiny

Common mullein

Corn Spurry

Creeping Beggarweed

Creeping Jenny

(creeping Charlie)

Cupid shaving brush

Cudweed

Daisy, English

Daisy, oxeye

Dandelion, common

Dandelion, false

Day flower

Deadnettle

Dichondra

Dock, broadleaf

Dock, curly

Dogbane

Dogfennel

Dollarweed (Pennywort)

Dove weed

Elderberry

Falseflax

False sunflower

Fiddleneck

Filaree, redstem

Filaree, whitestem

Fleabane, daisy (annual)

Fleabane, rough

Gallsoga, hairy

Gallsoga, smallflower

Goldenrod

Ground ivy

Groundsel

Gumweed

Hairy fleabane

Hawkweed, meadow

Hawkweed, mouseear

Hawkweed, orange

Hawkweed, yellow

Heal-all

Heartleaf drymary

Hemp

Henbit

Hoary cress

Horsenettle

Horseweed

Jimsonweed

Johnny-Jumpup violet

Knawel

Knotweed, prostrate

Kochia

Lambsquarters

Lespedeza

Mallow, alkali

Mallow, bristly

Mallow, common

Mallow, dwarf

Mallow, Venice

Marcela

Matchweed

Mexicanweed

Moneywort

Morningglory, bigroot

Morningglory, ivyleaf

Morningglory, Mexican

Morningglory, red

Morningglory, tall

Morningglory, woolly

Mugwort

Mustard, ball

Mustard, black

Mustard, blue

Mustard, hedge

Mustard, Indian

Mustard, tansy

Mustard, tumble

Mustard, white

Mustard, wormseed

Nettle, stinging (burning)

Nettle, tall

Nettle, wood

Parsley-piert

Parsnip

Pearlwort

Pennycress (frenchweed)

Pepperweed, perennial

Pepperweed, Virginia

(Peppergrass)

Pigweed, prostate

Pigweed, redroot

Pigweed, Russian

Pigweed, smooth

Pigweed, tumble

Pineapple weed

Plains coreopsis (tickseed)

Plantain, blackseed

Plantain, bracted

Plantain, broadleaf

Plantain, buckhorn

Plantain, hoary

Plantain, narrowleaf

Plantain, slender

Plantain, woolly

Poison ivy

Poison oak

Pokeweed

Prairie Sunflower

Prickly lettuce

(compass plant)

Prickly sida

Prostrate knotweed

Puncturevine

Purslane, common

Pusley, Brazil

Pusley, Florida

Pusley, large flower

Ragweed, bur

Ragweed, common

Ragweed, lanceleaf

Ragweed, Western

Redstem filaree

Rough cinquefoil

Scarlet pimpernel

Shepard's purse

Smartweed, ladythumb

Smartweed, pale

Smartweed, Pennsylvania

Smooth chaff-flower

Smooth dock

Sorrel, red (sheep)

Sowthistle, annual

Sowthistle, spiny

Spanish needles

Speedwell, bi-lobed

Speedwell, common

Speedwell, corn

Speedwell, ivyleaf

Speedwell, Persian

Speedwell, purslane

Speedwell, slender

Speedwell, snow

Speedwell, thymeleaf

Spiny amaranth

Spurge, spotted

Spurge, prostrate

Spurweed

Star of Bethlehem

Strawberry, India mock

Tansy ragwort

Tanweed

Thistle, blessed

Thistle, bull

Thistle, Canada

Thistle, Flodman's

(prairie)

Thistle, musk

Thistle, platte

Thistle, plumeless

Thistle, Russian

Thistle, Scotch

Thistle, wavyleaf

Thistle, yellow (horrible)

Trailing crownvetch

Velvetleaf

Vervain, blue

Vervain, hoary

Vervain, prostrate

Vervain, tall

Vetch, bird

Vetch, hairy

Vetch, milk

Vetch, narrowleaf

Vetch, wild

Virginia creeper

Western clematis

Western salsify

Wild aster

Wild buckwheat

Wild carrot

Wild four-o'clock

Wild garlic

Wild lettuce

Wild marigold

Wild mustard

(yellow rocket)

Wild onion

Wild parsnip

Wild radish

Wild rape

Wild strawberry

Wild sweet potato

Wild violet

Woodsorrel, creeping

Woodsorrel, violet

Woodsorrel, yellow

(Oxalis)

Woolly croton

Wormseed

Yarrow, common

Yarrow, Western

Yarrow, yellow

APPLICATION INSTRUCTIONS:

1. Mow lawn to normal height 1 to 2 days before application.

2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).

3. Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water before applying product.

4. Avoid applying when temperatures are consistently above 90°F.

5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

Spreader Settings

Spreader Directions:

Determining Your Lawn's Square Footage

LENGTH

WIDTH

To determine how much of this product to apply, you must determine the square footage of your lawn. An easy way to do this is to walk the length of your lawn, figuring that one pace equals about 3 feet. Do the same with the width of the lawn. Then multiply the length by the width to arrive at the total. Make sure you subtract the square footage of your home and driveway when calculating the total square footage of your lawn. If your lawn has an irregular shape, try dividing it up into a number of square areas. Calculate the square footage for each area, then add the figures together to reach your total lawn size.

Spreader Tips

The two main types of spreaders are the Drop Spreader, which drops the material straight down the width of the spreader as you push, and the Broadcast Spreader (also known as a rotary), which throws the material from the spreader.

| Broadcast Spreader | Setting 2.6 Lbs./1,000 Sq. Ft. | Drop Spreader | Setting 2.6 Lbs./1,000 Sq. Ft. |
|-------------------------|-----------------------------------|------------------|-----------------------------------|
| Agri-Fab | 4 | Earthway | 14 |
| Earthway EV-N Spred | 15 | Republic Drop | 4 |
| Republic EZGrow | 6½ | Scotts AccuGreen | 3½ |
| Scotts Deluxe Edgeguard | 4 | Sta-Green | 3½ |
| Scotts EasyGreen | 23 | | |
| Sta-Green | 3 | | |

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Broadcast Spreader:

Apply in a single pass by starting on the outside border and apply in a circular path working toward the center.

Drop Spreader:

First make two passes along the top of the area you'll be covering. Then make two passes along the bottom area. Then move back and forth along the long part of the lawn. On each pass, as you reach the strips at the top and bottom of the area to be covered, shut off the spreader. It's also important to shut off the spreader when you are stopped.

Note: if spreader model does not appear above please call 800-233-1067, 8 am - 5 pm ET, Monday – Friday for additional model settings.

The above spreader settings are approximate. To check your spreader: 1 lb. of material should cover approximately 385 sq. ft. (2.6 lbs. of material per 1,000 sq. ft.)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

Pesticide Disposal and Container Handling: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Questions or Comments? Call 1-800-233-1067.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

27-0-4

GUARANTEED ANALYSIS

Total Nitrogen (N) 27%

3.4% Ammoniacal Nitrogen

2.2% Water Insoluble Nitrogen*

18.0% Urea Nitrogen

3.4% Other Water Soluble Nitrogen*

Soluble Potash (K₂O) 4%

Sulfur (S) 3.8%

3.8% Combined Sulfur (S)

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.

* 5.6% Slowly Available Nitrogen from Methylene Ureas. F699

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-lebsea.com>.

YOUR SATISFACTION GUARANTEED

GREENVIEW® YARD WIDE GUARANTEE

OR YOUR MONEY BACK

Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem and simply send proof of purchase to Lebanon Seaboard Corporation, 1600 East Cumberland Street, Lebanon, PA 17042.

•For more information on our complete line of GreenView® Lawn and Garden Products, visit our website at www.greenviewfertilizer.com

•GreenView® is a registered trademark of Lebanon Seaboard Corporation.

•GREENSMART and Design is a trademark of Lebanon Seaboard Corporation.

Manufactured by:

Lebanon Seaboard Corporation

1600 East Cumberland Street

Lebanon, PA 17042

EPA Reg. No. 961-415

21-31169

i/24

(L) (D)

EPA Est. No. 961-PA-01, 961-IL-01

Superscript used is first letter of lot number.