

GreenView®

CRABGRASS CONTROL  
+LAWN FERTILIZER

GreenView®

CRABGRASS CONTROL +LAWN FERTILIZER

NOTICE: Read the entire label. Use only according to label directions.

Before buying this product, read the Warranty Section elsewhere on this label. If terms are not acceptable, return the unopened container at once.

For control of crabgrass and control or suppression of many other listed annual grass and broadleaf weeds in established lawns and ornamental turf.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Use Protective eyewear when using Battery Powered Spreader and Battery Powered Pail. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

**First Aid**

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

If inhaled:

Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.

• Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For Emergency Medical Treatment Information call (888)208-1368.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift or runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. DO NOT apply when weather conditions favor drift from treated areas. To protect the environment, DO NOT allow pesticide to enter or run off into storm drains, waterways, or surface waters. If you spill this product in calm weather, help to contain it, then and do not blow pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies and drainage systems.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

**How This Product Works**

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. GreenView Crabgrass Control Plus Lawn Fertilizer can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months of crabgrass control.

GreenView Crabgrass Control Plus Lawn Fertilizer provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only, but is effective up to 4 weeks after crabgrass has germinated. This generally corresponds to the time when crabgrass seedlings first become visible. DO NOT apply this product later than 4 weeks after crabgrass germination.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

Cool-Season Grasses	Warm-Season Grasses
Bentgrass, creeping	Bahiagrass
Bluegrass, Kentucky	Bermudagrass
Fescue, fine	Buffalograss
Fescue, tall	Carpetgrass
Ryegrass, perennial	Centipedegrass
	St. Augustinegrass
	Zoysiagrass

**Restrictions**

• DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.

• DO NOT allow people or pets to enter treated area until dust has settled.

• DO NOT apply more than 6.0 lbs. of this product per 1,000 sq. ft. per application and no more than 18.0 lbs. of this product/1,000 sq. ft. per year (equivalent to 0.5 lb. to 1.5 lbs. of active ingredient per acre, respectively).

• In the state of New York, DO NOT exceed 3.0 lbs. of this product per 1,000 sq. ft. per year (equivalent to 0.25 lb. of active ingredient per acre).

• DO NOT apply this product later than 4 weeks after crabgrass germination.

**Use Precautions**

GreenView Crabgrass Control Plus Lawn Fertilizer must be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, use the initial application of this product only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two more following seeding, sodding, or sprigging. Exception: In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied as the first seed.

• Use of this product on lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product to lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperature, or lack of moisture may result in turfgrass injury.

• For best results, cultural practices that disturb the soil, such as core, spike-, or hydroaeration, and verticulating, are to be done before application of this product. DO NOT apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

• Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application (splitting totaling 6.0 lbs./1,000 sq. ft. (0.5 lb. of active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When resowing, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

• DO NOT use this product later than 4 weeks after crabgrass germination.

• DO NOT apply this product